

The anomalous chiral perturbation theory meson Lagrangian to order p^6 revisited

T. Ebertshäuser,¹ H. W. Fearing,² S. Scherer¹

¹ *Institut für Kernphysik, Johannes Gutenberg-Universität, J. J. Becher-Weg 45, D-55099 Mainz, Germany*

² *TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia, Canada V6T 2A3*
(18 October 2001)

Abstract

We present a revised and extended construction of the mesonic Lagrangian density in chiral perturbation theory (ChPT) at order p^6 in the anomalous (or epsilon) sector, $\mathcal{L}_{6,\epsilon}$. After improving several aspects of the strategy we used originally, i.e., a more efficient application of partial integration, the implementation of so-called Bianchi identities, and additional trace relations, we find the new monomial sets to include 24 $SU(N_f)$, 23 $SU(3)$, and 5 $SU(2)$ elements. Furthermore, we introduce 8 supplementary terms due to the extension of the chiral group to $SU(N_f)_L \times SU(N_f)_R \times U(1)_V$.

12.39.Fe, 11.30.Rd

I. INTRODUCTION

Chiral perturbation theory for mesons [1–3] has proven to be a highly successful method for describing the interactions of the pseudoscalar octet (π, K, η) at low energies (for an overview of recent activities see, e.g., Ref. [4]). It is based on a chiral $SU(3)_L \times SU(3)_R$ symmetry of the QCD Lagrangian in the limit of massless u , d , and s quarks in combination with the assumption of a spontaneous symmetry breaking to $SU(3)_V$ in the groundstate. According to Goldstone’s theorem [5–7], each broken generator, i.e., each generator which does not annihilate the vacuum, gives rise to a massless Goldstone boson whose properties are tightly connected to the generator in question. In the present case, one expects eight pseudoscalar Goldstone bosons transforming as an octet under $SU(3)_V$ with vanishing interactions in the limit of zero energies. These Goldstone bosons are identified with the low-energy pseudoscalar octet, where the finite masses of the physical multiplet result from an explicit symmetry breaking due to the finite u , d , and s quark masses in the underlying QCD Lagrangian. As an analogous situation holds for $SU(2)$, we first consider the general case of $SU(N_f)$ and specialize to $N_f = 2$ and 3 later.

The effective Lagrangian density of ChPT is organized as a string of terms with an increasing number of covariant derivatives and quark mass terms,

$$\mathcal{L} = \mathcal{L}_2 + \mathcal{L}_4 + \mathcal{L}_6 + \cdots, \quad (1)$$

where the subscripts refer to the order in the momentum and quark mass expansion. The derivative expansion essentially takes account of the vanishing interaction at low energies with the explicit symmetry breaking being treated perturbatively. At each order the most general Lagrangian compatible with chiral symmetry, parity, and charge conjugation invariance is required. Here, we work in the framework of ordinary ChPT, where the quark mass term is counted as $\mathcal{O}(p^2)$, or, in other words, matrix elements are treated at a fixed ratio m_{quark}/p^2 [2]. For an overview of a different organization procedure we refer the reader to Ref. [8]. A systematic treatment of physical matrix elements is made possible by Weinberg’s power counting scheme [1]. For example, at $\mathcal{O}(p^4)$ one has to consider tree-level diagrams with exactly one vertex from $\mathcal{L}_4$ and an arbitrary number of vertices from $\mathcal{L}_2$ or one-loop diagrams with vertices from $\mathcal{L}_2$. The most general structure of $\mathcal{L}_4$ was first discussed by Gasser and Leutwyler [3] and contains 10 low-energy coupling constants L_i . Out of these, 8 are required to absorb infinities generated by one-loop diagrams from $\mathcal{L}_2$. The finite pieces are not predicted by chiral symmetry and have to be determined from experimental data.

The ChPT action functional, which is (apart from the special case of the Wess-Zumino-Witten term) the four-dimensional space-time integral over some chirally invariant Lagrangian density, generates at any given chiral order a finite-dimensional real vector space. The finite dimension is due to the fact that the basic building blocks can only be multiplied together in a finite number of different ways. Unfortunately, there seems to be neither a way to predict this dimension beforehand nor a general algorithm to decide whether a set of given structures is linearly independent or not. That is the reason why it is almost impossible to tell if a generating system is linearly independent so that it actually represents a basis of the above mentioned vector space. To our knowledge even the $\mathcal{L}_4$ of [3] has not formally been shown to be a basis, though it is without any doubt a generating system and countless calculations seem to confirm that the terms are independent. In Ref. [9] we systematically

wrote down the most general Lagrangian density of chiral order $\mathcal{O}(p^6)$, both for the normal ($\rightarrow 111$ $SU(3)$ terms) and anomalous ($\rightarrow 32$ $SU(3)$ terms) sector. Although these sets are also most likely to be generating systems, the normal sector was later shown to include redundant structures [10].

In the present paper, we provide a revised and slightly modified list of $SU(N_f)$ terms for the anomalous or epsilon sector, as well as its reduction to $SU(2)$ and $SU(3)$. For the sake of clarity, we repeat the analysis from the start, using a different notation which is closer to what has become the standard in this field. The initial ingredients and the general strategy are the same as used before. The main improvements in comparison to our former work are outlined one after the other in the following sections. In Sec. II, after defining different basic building blocks, which lead to nice simplifications of the partial-integration method (see Sec. III), we repeat the main features of our strategy. The so-called Bianchi identities for field strength tensors are introduced and implemented in Sec. IV. Section V contains a somewhat different approach to trace relations. In Sec. VI, additional structures triggered by the extension of the chiral group to $SU(N_f)_L \times SU(N_f)_R \times U(1)_V$ are worked out. A concluding comparison of the anomalous Lagrangian density to chiral order p^6 derived in Ref. [9] with the present work appears in Sec. VII. Section VIII gives a brief summary. Tables and eliminating relations are relegated to the Appendices.

II. MODIFIED BASIC BUILDING BLOCKS AND STRATEGY

In Ref. [9] a systematic construction of chiral Lagrangian densities was developed starting from structures of the type

$$[A]_{\pm} \equiv \frac{1}{2}(AU^{\dagger} \pm UA^{\dagger}), \quad (2)$$

where A is supposed to transform under the chiral group G as $A \xrightarrow{G} V_R A V_L^{\dagger}$. The partial-integration argument to be outlined in Sec. III may be simplified considerably by using a modified kind of basic building blocks

$$(A)_{\pm} \equiv 2u^{\dagger}[A]_{\pm}u, \quad (3)$$

which has already been used in a slightly different form by several authors (see, e.g., [10,11]). Here, u is defined via $u^2 \equiv U$. Note that $(A)_{\pm}$ obeys the transformation rule $(A)_{\pm} \xrightarrow{G} V(A)_{\pm}V^{\dagger}$, where $V \equiv \sqrt{V_R U V_L^{\dagger}}^{-1} V_R \sqrt{U}$ is the so-called compensator. The covariant derivative

$$\nabla_{\mu}(A)_{\pm} \equiv \partial_{\mu}(A)_{\pm} + [\Gamma_{\mu}, (A)_{\pm}] \quad (\text{N.B. : } \nabla_{\mu}(A)_{\pm} \xrightarrow{G} V \nabla_{\mu}(A)_{\pm} V^{\dagger}) \quad (4)$$

induced by the chiral connection $\Gamma_{\mu} = \frac{1}{2} [u^{\dagger}, \partial_{\mu}u] - \frac{i}{2} u^{\dagger} R_{\mu} u - \frac{i}{2} u L_{\mu} u^{\dagger}$ provides a key element for an easier and more efficient application of the total-derivative procedure, which is to be outlined in more detail shortly.

Since we always have to take single or multiple traces (denoted by $\langle \cdots \rangle$) to obtain Lagrangian density monomials, our final results will only differ by a number compared to the former structures

$$\langle (A_1)_\pm \dots (A_m)_\pm \rangle = 2^m \langle [A_1]_\pm \dots [A_m]_\pm \rangle. \quad (5)$$

Let us illustrate this fact by looking at the leading-order non-anomalous Lagrangian density

$$\mathcal{L}_2 = -\frac{F_0^2}{16} \langle (D_\mu U)_- (D^\mu U)_- \rangle + \frac{F_0^2}{4} \langle (\chi)_+ \rangle = -\frac{F_0^2}{4} \langle [D_\mu U]_- [D^\mu U]_- \rangle + \frac{F_0^2}{2} \langle [\chi]_+ \rangle.$$

As pointed out in Ref. [9], it is sufficient to restrict oneself to $[D^m U]_-$, $[D^n G]_+$, $[D^n H]_+$, and $[D^n \chi]_\pm$ (m, n integer with $m > 0, n \geq 0$). So, we simply have to substitute for those ingredients the new building blocks $(D^m U)_-$, $(D^n G)_+$, $(D^n H)_+$, and $(D^n \chi)_\pm$. Here we use the definitions $G^{\mu\nu} \equiv F_R^{\mu\nu} U + U F_L^{\mu\nu}$ and $H^{\mu\nu} \equiv F_R^{\mu\nu} U - U F_L^{\mu\nu}$, where $F_{L/R}^{\mu\nu}$ are evidently the field strength tensors belonging to the corresponding external fields R_μ and L_μ . The $\chi \equiv 2B_0(s + ip)$ block contains the external scalar and pseudoscalar fields, and B_0 is related to the scalar quark condensate $\langle 0 | \bar{q}q | 0 \rangle$ in the chiral limit. To simplify the comparison with other groups [9–11], we have given a translation prescription in Table I.

The so-called hierarchy strategy of Ref. [9] has proven to be extremely fruitful. Collecting $(D^n G_{\mu\nu})_+$, $(D^n H_{\mu\nu})_+$, and $(D^n \chi)_\pm$ for a moment in one symbol $(D^n \chi_{\mu\nu})_\pm$, one immediately finds that all possible terms at $\mathcal{O}(p^6)$ can either include no, one, two, or three $(D^n \chi_{\mu\nu})_\pm$ blocks which naturally defines four distinct sectors or levels to be considered. We say: the more $(D^n \chi_{\mu\nu})_\pm$ blocks are included in one monomial, the lower its level. Once all terms one can think of (at each level) are listed, one tries to generate relations to eliminate as many structures as possible. We always try to get rid of terms as high in the hierarchy as possible. In particular, with this strategy one ensures that the number of terms is minimal for the special case in which all external fields are set equal to zero.¹

Before explaining the improvements mentioned above in more detail, let us discuss the symmetry of multiple covariant derivatives. It turns out to be more useful not to assume the multiple covariant derivatives to be symmetrized from the very beginning. We implement the relations among unsymmetrized structures noting that just one general (i.e., non-contracted) index combination is actually independent in the sense of the hierarchy explained above. For double derivatives this statement reads

$$(D_\mu D_\nu A)_\pm = (D_\nu D_\mu A)_\pm + \frac{i}{4} [(A)_\pm, (G_{\mu\nu})_+] - \frac{i}{4} \{ (A)_\mp, (H_{\mu\nu})_+ \}. \quad (6)$$

Although there would not be any disadvantage in keeping these unsymmetrized monomials in our final set (N.B. : $(D_\mu D_\nu U)_-$ will be the only one to be kept), for aesthetical reasons we replace them by symmetrized ones making use of the formula

$$(D_\mu D_\nu U)_-^s \equiv \frac{1}{2} (\{D_\mu, D_\nu\} U)_- = (D_\mu D_\nu U)_- + \frac{i}{2} (H_{\mu\nu})_+. \quad (7)$$

¹The non-anomalous final $SU(N_f)$ set given in Ref. [10] is not minimal when setting all external fields to zero. In that case the structures Y_1 and Y_4 are not independent and can be eliminated. However, if we replace these two monomials by the terms (120) and (139) of Ref. [9], then the entire set will remain independent whether or not there are external fields and both structures, (120) and (139), vanish explicitly when the fields are set to zero.

We are now ready to initiate the construction procedure. First of all, we write down all conceivable (we find 74, see Table II) anomalous structures (satisfying P and C invariance, hermiticity, and chiral order p^6) in terms of our chirally invariant basic building blocks. We then collect as many relations as possible which follow from any of the mechanisms we are aware of. Those are

- (1) partial integration,
- (2) equation-of-motion argument,
- (3) epsilon relations,
- (4) Bianchi identities,
- (5) trace relations.

As detailed descriptions of methods (2) and (3) can be found in Ref. [9], we only comment on the new or improved items (1), (4), and (5). The resulting final set is shown in Table IV. There, the structures are again ordered according to the organization scheme introduced in IV.B of [9]. All explicit non-redundant relations are listed in Appendices A to E. There is no unique way to choose which of a set of monomials to keep. We basically follow the example of [9], i.e., in addition to the hierarchy approach we always try to get rid of multiple derivative terms, especially of the type $(D^2U)_-$. As emphasized above, and in Ref. [9], since one cannot prove the independence of the final set of monomials, there is no guarantee that another approach might not happen to find additional relations leading to a smaller final number. Yet we have not found an alternative way, giving rise to further reductions.

III. PARTIAL INTEGRATION (TOTAL-DERIVATIVE ARGUMENT)

Recalling the fact that a total derivative in the Lagrangian density does not change the equation of motion, we can generate relations of the following type

$$\begin{aligned} \partial_\mu \langle (A_1)_\pm \dots (A_m)_\pm \rangle + \overbrace{\langle [\Gamma_\mu, (A_1)_\pm \dots (A_m)_\pm] \rangle}^0 &= \langle \nabla_\mu [(A_1)_\pm \dots (A_m)_\pm] \rangle \\ &= \langle \nabla_\mu (A_1)_\pm \dots (A_m)_\pm \rangle + \dots + \langle (A_1)_\pm \dots \nabla_\mu (A_m)_\pm \rangle, \end{aligned} \quad (8)$$

where we made use of Eq. (4). This derivative shifting procedure is also valid for multiple traces. The advantage of our new basic building blocks stems from the relatively simple connection between the covariant derivative ∇_μ outside the block brackets and the covariant derivative D_μ inside

$$\nabla_\mu (A)_\pm = (D_\mu A)_\pm - \frac{1}{4} \{ (D_\mu U)_-, (A)_\mp \}. \quad (9)$$

Equation (9) is important because it helps to avoid extremely tedious algebraic manipulations one had to perform in the old framework of Ref. [9].²

² A combination of shifting derivatives back and forth and interchanging indices of multiple derivatives is sometimes referred to as index exchange.

Appendix A collects 23 non-redundant partial-integration relations (written in terms of the numbers indicated in Table II) which are used to eliminate dependent structures. Note that some of these relations might neglect contributions from lower levels. Since we always try to find the most general Lagrangian density at each level, in almost all cases those lower terms do not need to be known explicitly.

IV. BIANCHI IDENTITIES

As a consequence of the Jacobi identity

$$[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0, \quad (10)$$

certain combinations of covariant derivatives of the field strength tensors are not linearly independent. According to Eq. (18) of Ref. [9], the covariant derivative is defined as

$$D_\mu F_{\nu\rho}^R = \partial_\mu F_{\nu\rho}^R - i[R_\mu, F_{\nu\rho}^R]. \quad (11)$$

Now consider the linear combination

$$\begin{aligned} D_\mu F_{\nu\rho}^R + D_\nu F_{\rho\mu}^R + D_\rho F_{\mu\nu}^R &\equiv \sum_{\text{c.p.}\{\mu,\nu,\rho\}} D_\mu F_{\nu\rho}^R = \sum_{\text{c.p.}\{\mu,\nu,\rho\}} (\partial_\mu F_{\nu\rho}^R - i[R_\mu, F_{\nu\rho}^R]) \\ &= \sum_{\text{c.p.}\{\mu,\nu,\rho\}} \left(\partial_\mu \partial_\nu R_\rho - \partial_\mu \partial_\rho R_\nu - i[\partial_\mu R_\nu, R_\rho] \right. \\ &\quad \left. - i[R_\nu, \partial_\mu R_\rho] - i[R_\mu, \partial_\nu R_\rho - \partial_\rho R_\nu] - [R_\mu, [R_\nu, R_\rho]] \right) \\ &= 0, \end{aligned} \quad (12)$$

where use of the Schwarz theorem, relabelling of indices, and the Jacobi identity, Eq. (10), has been made. Observe that the cyclic permutation of the indices μ, ν , and ρ has been denoted by $\text{c.p.}\{\mu, \nu, \rho\}$. Repeating the same arguments for the independent field strength tensor $F_{\mu\nu}^L$, the additional new constraints can be summarized as

$$\sum_{\text{c.p.}\{\mu,\nu,\rho\}} D_\mu F_{\nu\rho}^{L/R} = 0, \quad (13)$$

which are referred to as the Bianchi identities (see, e.g., Refs. [12,13]). Equation (13) does not require that $F_{\mu\nu}^{R/L}$ satisfy any equations of motion. In terms of our building blocks $(D_\mu U)_-$, $(G_{\mu\nu})_+$, and $(H_{\mu\nu})_+$ the Bianchi identities read

$$\sum_{\text{c.p.}\{\mu,\nu,\rho\}} (D_\mu G_{\nu\rho})_+ = -\frac{1}{4} \sum_{\text{c.p.}\{\mu,\nu,\rho\}} [(D_\mu U)_-, (H_{\nu\rho})_+], \quad (14)$$

$$\sum_{\text{c.p.}\{\mu,\nu,\rho\}} (D_\mu H_{\nu\rho})_+ = -\frac{1}{4} \sum_{\text{c.p.}\{\mu,\nu,\rho\}} [(D_\mu U)_-, (G_{\nu\rho})_+]. \quad (15)$$

We stress that in Ref. [9] each term (though in the old notation) of the sum on the left-hand side of Eqs. (14) and (15) was treated as an independent element.

In Appendix D one finds 7 eliminating relations induced by Bianchi identities. At this point we need to stress that sometimes a structure can be eliminated due to several arguments. Therefore, the number of non-redundant relations belonging to a specific subprocedure can vary, while the total number of all relations is constant.

V. TRACE RELATIONS

As all the structures one can think of either involve single or multiple traces, one is particularly interested in finding relations between those traces. We know from the Cayley-Hamilton theorem that any $n \times n$ matrix A is a solution of its associated characteristic polynomial χ_A . For $n = 2$ this statement reads

$$0 = \chi_A(A) = A^2 - \langle A \rangle A + \det(A)1 = A^2 - \langle A \rangle A + \frac{1}{2}(\langle A \rangle^2 - \langle A^2 \rangle)1. \quad (16)$$

Setting $A = A_1 + A_2$ in (16) and making use of $\chi_{A_1}(A_1) = 0 = \chi_{A_2}(A_2)$ one ends up with the matrix equation

$$F_2(A_1, A_2) \equiv \{A_1, A_2\} - \langle A_1 \rangle A_2 - \langle A_2 \rangle A_1 + \langle A_1 \rangle \langle A_2 \rangle 1 - \langle A_1 A_2 \rangle 1 = 0 \quad (17)$$

which is the central piece of information needed to derive the trace relations. The analogous $n = 3$ equation is slightly more complex

$$\begin{aligned} F_3(A_1, A_2, A_3) \equiv & A_1\{A_2, A_3\} + A_2\{A_3, A_1\} + A_3\{A_1, A_2\} \\ & - \langle A_1 \rangle \{A_2, A_3\} - \langle A_2 \rangle \{A_3, A_1\} - \langle A_3 \rangle \{A_1, A_2\} \\ & + \langle A_1 \rangle \langle A_2 \rangle A_3 + \langle A_2 \rangle \langle A_3 \rangle A_1 + \langle A_3 \rangle \langle A_1 \rangle A_2 \\ & - \langle A_1 A_2 \rangle A_3 - \langle A_3 A_1 \rangle A_2 - \langle A_2 A_3 \rangle A_1 \\ & - \langle A_1 A_2 A_3 \rangle 1 - \langle A_1 A_3 A_2 \rangle 1 \\ & + \langle A_1 A_2 \rangle \langle A_3 \rangle 1 + \langle A_3 A_1 \rangle \langle A_2 \rangle 1 + \langle A_2 A_3 \rangle \langle A_1 \rangle 1 \\ & - \langle A_1 \rangle \langle A_2 \rangle \langle A_3 \rangle 1 = 0. \end{aligned} \quad (18)$$

We can now derive trace relations by simply multiplying Eq. (17) or Eq. (18) with another arbitrary matrix of adequate dimension and finally taking the trace of the whole construction, i.e.,

$$0 = \langle F_2(A_1, A_2) A_3 \rangle, \quad (19a)$$

$$0 = \langle F_3(A_1, A_2, A_3) A_4 \rangle. \quad (19b)$$

Note that A_i may be any $n \times n$ matrix, even a string of our basic building blocks. Although we have never come across a trace relation that could not be obtained in the manner explained above, we are not aware of a general proof showing that any kind of trace relation must be related to the Cayley-Hamilton theorem.

Equation (19b) is actually identical to Eq. (A7) of Ref. [9]. In Appendix E we tabulate the 19 $SU(2)$ relations which follow from Eq. (19a) plus a single $SU(3)$ relation which was missed in Ref. [9].

VI. EXTENSION OF THE CHIRAL GROUP

So far we have presented the most general anomalous $SU(N_f)_L \times SU(N_f)_R$ invariant Lagrangian density (respecting external fields v_μ, a_μ, s , and p) at chiral order p^6 from which one can extract the $SU(2)$ or $SU(3)$ version by applying the respective trace relations. Since

$(G_{\mu\nu})_+$ and $(H_{\mu\nu})_+$ are both supposed to be traceless and the $SU(2)$ quark charge matrix $Q = \text{diag}(\frac{2}{3}, -\frac{1}{3})$ is not, even the special case of electromagnetism ($L_\mu = R_\mu = -eA_\mu Q$) is not fully included in the general $SU(2)$ formulae. As Q happens to be traceless in $SU(3)$, there is no such problem in this case. If we want to do away with this apparent weakness, we need to extend the chiral group to $SU(N_f)_L \times SU(N_f)_R \times U(1)_V$. For $N_f = 2$ the electromagnetic interaction is then fully included by setting $L_\mu = R_\mu = -\frac{e}{2}A_\mu\tau_3$ and $v_\mu^{(s)} = -\frac{e}{2}A_\mu$, where $v_\mu^{(s)}$ is the $U(1)_V$ gauge field. Equivalently, one might argue that the traceless fields L_μ and R_μ have to be replaced by non-traceless ones of the form $\tilde{L}_\mu = L_\mu + \frac{1}{3}v_\mu^{(s)}1_{N_f \times N_f}$ and $\tilde{R}_\mu = R_\mu + \frac{1}{3}v_\mu^{(s)}1_{N_f \times N_f}$, respectively.

The extension basically comes down to the fact that we get one more independent elementary building block of chiral order p^2 , namely the field strength tensor $v_{\mu\nu}^{(s)} = \partial_\mu v_\nu^{(s)} - \partial_\nu v_\mu^{(s)}$. Again, we write down all (30) anomalous p^6 structures including at least one $v_{\mu\nu}^{(s)}$ (see Table III). Observe that at lower orders there is no way to construct such terms. We then use the mechanisms of the previous sections to eliminate as many of the additional monomials as possible. Since trace relations do not induce reductions in this field, our result is valid for any N_f , although we are primarily interested in $N_f = 2$. While all explicit relations can be found in the Appendices, the final set of additional structures is shown in Table VI.

VII. COMPARISON WITH OUR FORMER SET

According to the present analysis the most general anomalous $SU(N_f)$ Lagrangian density of chiral order $\mathcal{O}(p^6)$, $\mathcal{L}_{6,\epsilon}$, must not include more than 24 elements. We have started with 74 initial $SU(N_f)$ structures and then worked out 50 eliminating relations (trace relations excluded). We have tried to keep as many of the 32 structures of Ref. [9] (up to signs and factors) as possible. However, some terms listed therein, such as A_3 and A_{24} , are unnecessarily complicated. Therefore, it turns out to be more appropriate to respectively replace these elements by $\mathbf{L}_{19}^{6,\epsilon}$ and $\mathbf{L}_{20}^{6,\epsilon}$ which is justified by the total-derivative argument of Sec. III.

The necessary reduction from 32 to 24 structures is then obtained in the following way. First of all, using once more the improved partial-integration method, A_{10} and A_{11} can be shown to be redundant. The application of the Bianchi identities implies the further elimination of $A_1, A_2, A_9, A_{17}, A_{18}$, and A_{23} . The exact correspondence between the new low-energy constants $\mathbf{L}_i^{6,\epsilon}$ and the old A_j is shown in Table VII.

Finally, one $SU(3)$ trace relation can be implemented to get rid of A_{30} .

VIII. SUMMARY

In Ref. [9] we presented a list of 32 anomalous $SU(N_f)$ (the same number was valid for $SU(3)$) monomials of chiral order $\mathcal{O}(p^6)$ which constitute the most general mesonic ChPT Lagrangian density $\mathcal{L}_{6,\epsilon}$. After carefully revisiting the entire project, our list has been found to be redundant. Thanks to a more efficient use of partial integration, the implementation of so-called Bianchi identities for field strength tensors, and an additional application of a trace relation we have ended up with 24 $SU(N_f)$, 23 $SU(3)$, and 5 $SU(2)$ slightly modified elements. Our final sets are thus considerably smaller than those proposed

by [11,14], which are both incomplete and redundant. Furthermore, we have constructed 8 additional structures which arise due to the extension of the chiral group to $SU(N_f)_L \times SU(N_f)_R \times U(1)_V$. Besides the general interest, the latter group needs to be considered when consistently treating electromagnetic reactions in the $SU(2)$ framework.

ACKNOWLEDGMENTS

We would like to thank J. Bijnens and G. Colangelo for drawing our attention to the Bianchi identities. This work was supported in part by the Deutsche Forschungsgemeinschaft (SFB 443) and in part by the Natural Sciences and Engineering Research Council of Canada. T. E. would like to thank the theory group at TRIUMF for its hospitality and the German Academic Exchange Service (DAAD) for financial support.

TABLES

our notation	Ref. [9]	Ref. [10]	Ref. [11]
$(\chi)_\pm$	$2u^\dagger[\chi]_\pm u$	$\chi_\pm$	$2M^\pm$
$(D_\mu U)_-$	$2u^\dagger[D_\mu U]_- u$	$-2iu_\mu$	$4\Delta_\mu$
$(G^{\mu\nu})_+$	$2u^\dagger[G^{\mu\nu}]_+ u$	$2f_+^{\mu\nu}$	$-2iF^{+\mu\nu}$
$(H^{\mu\nu})_+$	$2u^\dagger[H^{\mu\nu}]_+ u$	$-2f_-^{\mu\nu}$	$2iF^{-\mu\nu}$

TABLE I. Translation prescription.

#	Initial Structures
1	$i \langle (D_\lambda D^\lambda U)_- (D_\mu U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
2	$i \langle (D_\lambda D_\mu U)_- \{ (D^\lambda U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (D_\nu U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
3	$i \langle (D_\lambda D_\mu U)_- \{ (D_\nu U)_- (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (D^\lambda U)_- (D_\nu U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
4	$i \langle (\chi)_- (D_\mu U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
5	$\langle (D^\lambda G_{\lambda\mu})_+ (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
6	$\langle (D_\lambda G_{\mu\nu})_+ \{ (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
7	$\langle (D_\lambda G_{\mu\nu})_+ (D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
8	$\langle (D_\mu G_{\lambda\nu})_+ \{ (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
9	$\langle (D_\mu G_{\lambda\nu})_+ (D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
10	$\langle (D_\mu G_{\nu\alpha})_+ \{ (D_\lambda U)_- (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- (D_\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
11	$\langle (G_{\mu\nu})_+ \{ (D_\lambda D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (D_\lambda D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
12	$\langle (G_{\lambda\mu})_+ \{ (D^\lambda D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (D^\lambda D_\nu U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
13	$\langle (G_{\mu\nu})_+ \{ (D_\lambda D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- (D_\lambda D_\alpha U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
14	$\langle (G_{\mu\nu})_+ \{ (D_\lambda D_\alpha U)_- (D_\beta U)_- (D^\lambda U)_- - (D^\lambda U)_- (D_\beta U)_- (D_\lambda D_\alpha U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
15	$\langle (G_{\mu\nu})_+ (D_\alpha U)_- (D_\lambda D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
16	$\langle (G_{\lambda\mu})_+ (D_\nu U)_- (D^\lambda D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
17	$\langle (G_{\mu\nu})_+ \{ (D_\lambda U)_- (D^\lambda D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda D_\alpha U)_- (D_\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
18	$\langle (D_\lambda D_\mu H_{\nu\alpha})_+ \{ (D^\lambda U)_- (D_\beta U)_- + (D_\beta U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
19	$\langle (D_\lambda H_{\mu\nu})_+ \{ (D^\lambda D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D^\lambda D_\alpha U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
20	$\langle (D_\mu H_{\lambda\nu})_+ \{ (D^\lambda D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D^\lambda D_\alpha U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
21	$\langle (D_\mu H_{\nu\alpha})_+ \{ (D_\lambda D^\lambda U)_- (D_\beta U)_- + (D_\beta U)_- (D_\lambda D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
22	$\langle (D_\mu H_{\nu\alpha})_+ \{ (D_\lambda D_\beta U)_- (D^\lambda U)_- + (D^\lambda U)_- (D_\lambda D_\beta U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
23	$\langle (H_{\mu\nu})_+ \{ (D_\lambda D^\lambda D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\lambda D^\lambda D_\alpha U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
24	$\langle (H_{\lambda\mu})_+ \{ (D^\lambda U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (D_\nu U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
25	$\langle (H_{\lambda\mu})_+ \{ (D_\nu U)_- (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (D^\lambda U)_- (D_\nu U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
26	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (D^\lambda U)_- (D_\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
27	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- + (D_\beta U)_- (D^\lambda U)_- (D_\alpha U)_- (D_\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
28	$\langle (D_\lambda D_\mu H_{\nu\alpha})_+ \rangle \langle (D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
29	$\langle (D_\lambda D^\lambda D_\mu U)_- \rangle \langle (H_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
30	$\langle (H_{\mu\nu})_+ (D_\alpha U)_- \rangle \langle (D_\beta U)_- (D_\lambda U)_- (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
31	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D_\alpha U)_- + (D_\alpha U)_- (D_\lambda U)_- \} \rangle \langle (D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
32	$\langle (D_\mu \chi)_+ (D_\nu H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
33	$\langle (D_\mu \chi)_+ \{ (G_{\nu\alpha})_+ (D_\beta U)_- - (D_\beta U)_- (G_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
34	$\langle (D_\mu \chi)_- \{ (H_{\nu\alpha})_+ (D_\beta U)_- + (D_\beta U)_- (H_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$

#	continuation
35	$\langle (D_\mu G_{\nu\alpha})_+ \{(\chi)_+ (D_\beta U)_- - (D_\beta U)_- (\chi)_+\} \rangle \epsilon^{\mu\nu\alpha\beta}$
36	$\langle (D_\mu H_{\nu\alpha})_+ \{(\chi)_- (D_\beta U)_- + (D_\beta U)_- (\chi)_-\} \rangle \epsilon^{\mu\nu\alpha\beta}$
37	$i \langle (D^\lambda G_{\lambda\mu})_+ \{ (G_{\nu\alpha})_+ (D_\beta U)_- + (D_\beta U)_- (G_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
38	$i \langle (D^\lambda G_{\mu\nu})_+ \{ (G_{\lambda\alpha})_+ (D_\beta U)_- + (D_\beta U)_- (G_{\lambda\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
39	$i \langle (D_\lambda G_{\mu\nu})_+ \{ (G_{\alpha\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (G_{\alpha\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
40	$i \langle (D_\mu G_{\lambda\nu})_+ \{ (G^\lambda_\alpha)_+ (D_\beta U)_- + (D_\beta U)_- (G^\lambda_\alpha)_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
41	$i \langle (D_\mu G_{\lambda\nu})_+ \{ (G_{\alpha\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (G_{\alpha\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
42	$i \langle (D_\mu G_{\nu\alpha})_+ \{ (G_{\lambda\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (G_{\lambda\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
43	$i \langle (D^\lambda H_{\lambda\mu})_+ \{ (H_{\nu\alpha})_+ (D_\beta U)_- + (D_\beta U)_- (H_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
44	$i \langle (D^\lambda H_{\mu\nu})_+ \{ (H_{\lambda\alpha})_+ (D_\beta U)_- + (D_\beta U)_- (H_{\lambda\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
45	$i \langle (D_\lambda H_{\mu\nu})_+ \{ (H_{\alpha\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (H_{\alpha\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
46	$i \langle (D_\mu H_{\lambda\nu})_+ \{ (H^\lambda_\alpha)_+ (D_\beta U)_- + (D_\beta U)_- (H^\lambda_\alpha)_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
47	$i \langle (D_\mu H_{\lambda\nu})_+ \{ (H_{\alpha\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (H_{\alpha\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
48	$i \langle (D_\mu H_{\nu\alpha})_+ \{ (H_{\lambda\beta})_+ (D^\lambda U)_- + (D^\lambda U)_- (H_{\lambda\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
49	$i \langle (D_\lambda D^\lambda U)_- (G_{\mu\nu})_+ (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
50	$i \langle (D^\lambda D_\mu U)_- \{ (G_{\lambda\nu})_+ (G_{\alpha\beta})_+ + (G_{\alpha\beta})_+ (G_{\lambda\nu})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
51	$i \langle (D_\lambda D^\lambda U)_- (H_{\mu\nu})_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
52	$i \langle (D^\lambda D_\mu U)_- \{ (H_{\lambda\nu})_+ (H_{\alpha\beta})_+ + (H_{\alpha\beta})_+ (H_{\lambda\nu})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
53	$\langle (\chi)_- \{ (G_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (G_{\mu\nu})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
54	$\langle (\chi)_+ \{ (H_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- + (D_\beta U)_- (D_\alpha U)_- (H_{\mu\nu})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
55	$i \langle (G_{\lambda\mu})_+ \{ (H^\lambda_\nu)_+ (D_\alpha U)_- (D_\beta U)_- - (D_\beta U)_- (D_\alpha U)_- (H^\lambda_\nu)_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
56	$i \langle (G_{\lambda\mu})_+ \{ (H_{\nu\alpha})_+ (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- (H_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
57	$i \langle (G_{\lambda\mu})_+ \{ (H_{\nu\alpha})_+ (D_\beta U)_- (D^\lambda U)_- - (D^\lambda U)_- (D_\beta U)_- (H_{\nu\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
58	$i \langle (G_{\mu\nu})_+ \{ (H_{\lambda\alpha})_+ (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- (H_{\lambda\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
59	$i \langle (G_{\mu\nu})_+ \{ (H_{\lambda\alpha})_+ (D_\beta U)_- (D^\lambda U)_- - (D^\lambda U)_- (D_\beta U)_- (H_{\lambda\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
60	$i \langle (G_{\mu\nu})_+ \{ (H_{\alpha\beta})_+ (D_\lambda U)_- (D^\lambda U)_- - (D^\lambda U)_- (D_\lambda U)_- (H_{\alpha\beta})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
61	$\langle (\chi)_- (D_\mu U)_- (G_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
62	$i \langle (G_{\lambda\mu})_+ \{ (D^\lambda U)_- (H_{\nu\alpha})_+ (D_\beta U)_- - (D_\beta U)_- (H_{\nu\alpha})_+ (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
63	$i \langle (G_{\lambda\mu})_+ (D_\nu U)_- (H^\lambda_\alpha)_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
64	$i \langle (G_{\mu\nu})_+ \{ (D^\lambda U)_- (H_{\lambda\alpha})_+ (D_\beta U)_- - (D_\beta U)_- (H_{\lambda\alpha})_+ (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
65	$\langle (D_\mu \chi)_- \rangle \langle (H_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
66	$\langle (\chi)_- \rangle \langle (D_\mu H_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
67	$\langle (\chi)_- \rangle \langle (G_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
68	$\langle (\chi)_+ (D_\mu U)_- \rangle \langle (H_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
69	$i \langle (\chi)_+ \{ (G_{\mu\nu})_+ (H_{\alpha\beta})_+ - (H_{\alpha\beta})_+ (G_{\mu\nu})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
70	$i \langle (\chi)_- (G_{\mu\nu})_+ (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
71	$i \langle (\chi)_- (H_{\mu\nu})_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
72	$\langle (G_{\mu\nu})_+ \{ (G_{\lambda\alpha})_+ (H^\lambda_\beta)_+ + (H^\lambda_\beta)_+ (G_{\lambda\alpha})_+ \} \rangle \epsilon^{\mu\nu\alpha\beta}$
73	$i \langle (\chi)_- \rangle \langle (G_{\mu\nu})_+ (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
74	$i \langle (\chi)_- \rangle \langle (H_{\mu\nu})_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$

TABLE II. List of 74 anomalous $SU(N_f)$ structures of order $\mathcal{O}(p^6)$ we start with. Besides the hierarchy argument discussed in the text (the four different categories are separated by simple lines) our ordering scheme is supposed to respect the rule: single traces $>$ multiple traces.

#	Initial Structures
1'	$\partial^\lambda v_{\lambda\mu}^{(s)} \langle (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
2'	$\partial_\lambda v_{\mu\nu}^{(s)} \langle (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
3'	$\partial_\mu v_{\lambda\nu}^{(s)} \langle (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
4'	$v_{\lambda\mu}^{(s)} \langle (D^\lambda D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
5'	$v_{\mu\nu}^{(s)} \langle (D_\lambda D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
6'	$v_{\mu\nu}^{(s)} \langle (D_\lambda D_\alpha U)_- \{ (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
7'	$i \partial^\lambda v_{\lambda\mu}^{(s)} \langle (G_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
8'	$i \partial^\lambda v_{\mu\nu}^{(s)} \langle (G_{\lambda\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
9'	$i \partial_\lambda v_{\mu\nu}^{(s)} \langle (G_{\alpha\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
10'	$i \partial_\mu v_{\lambda\nu}^{(s)} \langle (G^\lambda_\alpha)_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
11'	$i \partial_\mu v_{\lambda\nu}^{(s)} \langle (G_{\alpha\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
12'	$i \partial_\mu v_{\nu\alpha}^{(s)} \langle (G_{\lambda\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
13'	$i v_{\mu\nu}^{(s)} \langle (D^\lambda G_{\lambda\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
14'	$i v_{\lambda\mu}^{(s)} \langle (D^\lambda G_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
15'	$i v_{\mu\nu}^{(s)} \langle (D_\lambda G_{\alpha\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
16'	$i v_{\lambda\mu}^{(s)} \langle (D_\nu G^\lambda_\alpha)_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
17'	$i v_{\mu\nu}^{(s)} \langle (D_\alpha G_{\lambda\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
18'	$i v_{\lambda\mu}^{(s)} \langle (D_\nu G_{\alpha\beta})_+ (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
19'	$i v_{\mu\nu}^{(s)} \langle (G_{\alpha\beta})_+ (D_\lambda D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
20'	$i v_{\lambda\mu}^{(s)} \langle (G_{\nu\alpha})_+ (D^\lambda D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
21'	$i v_{\mu\nu}^{(s)} \langle (G_{\lambda\alpha})_+ (D^\lambda D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
22'	$i v_{\lambda\mu}^{(s)} \langle (H^\lambda_\nu)_+ (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
23'	$i v_{\lambda\mu}^{(s)} \langle (H_{\nu\alpha})_+ \{ (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
24'	$i v_{\mu\nu}^{(s)} \langle (H_{\lambda\alpha})_+ \{ (D^\lambda U)_- (D_\beta U)_- - (D_\beta U)_- (D^\lambda U)_- \} \rangle \epsilon^{\mu\nu\alpha\beta}$
25'	$v_{\mu\nu}^{(s)} \langle (\chi)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
26'	$i v_{\mu\nu}^{(s)} \langle (\chi)_- (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
27'	$v_{\lambda\mu}^{(s)} \langle (G^\lambda_\nu)_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
28'	$v_{\lambda\mu}^{(s)} \langle (G_{\nu\alpha})_+ (H^\lambda_\beta)_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
29'	$v_{\mu\nu}^{(s)} \langle (G_{\lambda\alpha})_+ (H^\lambda_\beta)_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
30'	$i v_{\mu\nu}^{(s)} v_{\alpha\beta}^{(s)} \langle (\chi)_- \rangle \epsilon^{\mu\nu\alpha\beta}$

TABLE III. List of 30 additional anomalous p^6 structures due to the extension of the chiral group.

$\mathbf{L}_i^{6,\epsilon}$	#	$SU(N_f)$	$SU(3)$	$SU(2)$
3	69	$i \langle (\chi)_+ \{ (G_{\mu\nu})_+ (H_{\alpha\beta})_+ - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	$\times$
8	70	$i \langle (\chi)_- (G_{\mu\nu})_+ (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	$\times$
9	73	$i \langle (\chi)_- \rangle \langle (G_{\mu\nu})_+ (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
19	37	$i \langle (D^\lambda G_{\lambda\mu})_+ \{ (G_{\nu\alpha})_+ (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
1	54	$\langle (\chi)_+ \{ (H_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	$\times$
2	68	$\langle (\chi)_+ (D_\mu U)_- \rangle \langle (D_\nu U)_- (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
5	53	$\langle (\chi)_- \{ (G_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
6	61	$\langle (\chi)_- (D_\mu U)_- (G_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	$\times$
7	67	$\langle (\chi)_- \rangle \langle (G_{\mu\nu})_+ (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
13	14 ^s	$\langle (G_{\mu\nu})_+ \{ (D^\lambda D_\alpha U)_- (D_\beta U)_- (D_\lambda U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
14	13 ^s	$\langle (G_{\mu\nu})_+ \{ (D_\lambda D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
10	71	$i \langle (\chi)_- (H_{\mu\nu})_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	$\times$
11	74	$i \langle (\chi)_- \rangle \langle (H_{\mu\nu})_+ (H_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
20	43	$i \langle (D^\lambda H_{\lambda\mu})_+ \{ (H_{\nu\alpha})_+ (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
21	55	$i \langle (G_{\lambda\mu})_+ \{ (H^\lambda_\nu)_+ (D_\alpha U)_- (D_\beta U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
22	57	$i \langle (G_{\lambda\mu})_+ \{ (H_{\nu\alpha})_+ (D_\beta U)_- (D^\lambda U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
23	60	$i \langle (G_{\mu\nu})_+ \{ (H_{\alpha\beta})_+ (D_\lambda U)_- (D^\lambda U)_- - \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
24	63	$i \langle (G_{\lambda\mu})_+ (D_\nu U)_- (H^\lambda_\alpha)_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
4	4	$i \langle (\chi)_- (D_\mu U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
12	2 ^s	$i \langle (D_\lambda D_\mu U)_-^s \{ (D^\lambda U)_- (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
15	27	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D_\alpha U)_- (D^\lambda U)_- (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	—	—
16	26	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D^\lambda U)_- (D_\alpha U)_- (D_\beta U)_- + \text{rev} \} \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
17	31	$\langle (H_{\mu\nu})_+ \{ (D_\lambda U)_- (D_\alpha U)_- + \text{rev} \} \rangle \langle (D^\lambda U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—
18	30	$\langle (H_{\mu\nu})_+ (D_\alpha U)_- \rangle \langle (D_\beta U)_- (D_\lambda U)_- (D^\lambda U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$	$\times$	—

TABLE IV. The final $SU(N_f)$ set. The remaining structures and their associated low-energy constants (LECs) have been relabelled: terms with a $(\chi)_\pm$ block from $\mathbf{L}_{11}^{6,\epsilon}$ to $\mathbf{L}_{11}^{6,\epsilon}$ and those without from $\mathbf{L}_{12}^{6,\epsilon}$ to $\mathbf{L}_{24}^{6,\epsilon}$. Afterwards they have been classified according to their leading-order expansion in terms of Goldstone boson and electromagnetic fields, see Table V. The abbreviation “rev” stands for the reverse order. Trace relations lead to the corresponding $SU(3)$ and $SU(2)$ sets, which are indicated by a $\times$ in the respective columns. Recall that under the traces the new $(\dots)_\pm$ blocks are equivalent to the old $[\dots]_\pm$ up to a factor 2.

reaction type	LECs
$\phi + 2\gamma^*$	3, 8, 9, 19
$3\phi + \gamma^*$	1, 2, 5, 6, 7, 13, 14
$3\phi + 2\gamma^*$	10, 11, 20, 21, 22, 23, 24
5ϕ	4, 12
$5\phi + \gamma^*$	15, 16, 17, 18

TABLE V. Classification of LECs according to their leading-order contribution allowing at most electromagnetic external fields; the photons may be either virtual or real. This classification defines the ordering scheme in Table IV.

$\mathbf{L}_i^{6,\epsilon}$	#	additional structures
2'	26'	$i v_{\mu\nu}^{(s)} \langle (\chi)_- (G_{\alpha\beta})_+ \rangle \epsilon^{\mu\nu\alpha\beta}$
3'	30'	$i v_{\mu\nu}^{(s)} v_{\alpha\beta}^{(s)} \langle (\chi)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
5'	13'	$i v_{\mu\nu}^{(s)} \langle (D^\lambda G_{\lambda\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
6'	7'	$i \partial^\lambda v_{\lambda\mu}^{(s)} \langle (G_{\nu\alpha})_+ (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
1'	25'	$v_{\mu\nu}^{(s)} \langle (\chi)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
4'	1'	$\partial^\lambda v_{\lambda\mu}^{(s)} \langle (D_\nu U)_- (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
7'	22'	$i v_{\lambda\mu}^{(s)} \langle (H^\lambda{}_\nu)_+ (D_\alpha U)_- (D_\beta U)_- \rangle \epsilon^{\mu\nu\alpha\beta}$
8'	29'	$v_{\mu\nu}^{(s)} \langle (G_{\lambda\alpha})_+ (H^\lambda{}_\beta)_+ \rangle \epsilon^{\mu\nu\alpha\beta}$

TABLE VI. Final set of additional structures valid for arbitrary N_f .

LECs	$\mathbf{L}_1^{6,\epsilon}$	$\mathbf{L}_2^{6,\epsilon}$	$\mathbf{L}_3^{6,\epsilon}$	$\mathbf{L}_4^{6,\epsilon}$	$\mathbf{L}_5^{6,\epsilon}$	$\mathbf{L}_6^{6,\epsilon}$	$\mathbf{L}_7^{6,\epsilon}$	$\mathbf{L}_8^{6,\epsilon}$
Ref. [9]	$16A_{15}$	$-16A_{16}$	$-8A_5$	$16A_{28}$	$16A_{12}$	$-16A_{13}$	$16A_{14}$	$8A_4$
LECs	$\mathbf{L}_9^{6,\epsilon}$	$\mathbf{L}_{10}^{6,\epsilon}$	$\mathbf{L}_{11}^{6,\epsilon}$	$\mathbf{L}_{12}^{6,\epsilon}$	$\mathbf{L}_{13}^{6,\epsilon}$	$\mathbf{L}_{14}^{6,\epsilon}$	$\mathbf{L}_{15}^{6,\epsilon}$	$\mathbf{L}_{16}^{6,\epsilon}$
Ref. [9]	$8A_6$	$8A_{25}$	$8A_{26}$	$32A_{27}$	$16A_8$	$16A_7$	$-32A_{30}$	$-32A_{29}$
LECs	$\mathbf{L}_{17}^{6,\epsilon}$	$\mathbf{L}_{18}^{6,\epsilon}$	$\mathbf{L}_{19}^{6,\epsilon}$	$\mathbf{L}_{20}^{6,\epsilon}$	$\mathbf{L}_{21}^{6,\epsilon}$	$\mathbf{L}_{22}^{6,\epsilon}$	$\mathbf{L}_{23}^{6,\epsilon}$	$\mathbf{L}_{24}^{6,\epsilon}$
Ref. [9]	$32A_{32}$	$32A_{31}$	n.d.c.	n.d.c.	$16A_{20}$	$-16A_{22}$	$16A_{19}$	$16A_{21}$

TABLE VII. Numerical correspondence between the $SU(N_f)$ low-energy constants (LECs) defined in Table IV and those of our former set. Observe that “n.d.c.” stands for “no direct correspondence.”

APPENDIX A: RELATIONS INDUCED BY PARTIAL INTEGRATION

$$\begin{aligned}
(1) - (2) + (3) &= 0, \\
(5) + (12) + (16) &= 0, \\
(6) + (11) + (14) + (17) &= 0, \\
(7) + (13) + (15) &= 0, \\
(8) - (12) &= 0, \\
(9) + (16) &= 0, \\
(10) + (13) + (17) &= 0, \\
(18) + (21) + (22) + \frac{1}{16}(26) + \frac{1}{8}(27) &= 0, \\
(20) - \frac{1}{16}(24) - \frac{1}{16}(25) &= 0, \\
(22) + \frac{1}{16}(26) &= 0, \\
(19) + (23) - \frac{1}{16}(26) - \frac{1}{8}(27) &= 0, \\
(21) + (23) - \frac{1}{8}(26) &= 0, \\
(28) - \frac{1}{8}(31) &= 0, \\
(29) + \frac{1}{4}(30) &= 0, \\
(32) + \frac{1}{4}(36) + \frac{1}{16}(54) &= 0, \\
(32) - \frac{1}{4}(34) &= 0, \\
(37) + (38) - (50) &= 0, \\
(39) + (49) &= 0, \\
(41) + (42) + (50) &= 0, \\
(43) + (44) - (52) &= 0, \\
(45) + (51) &= 0, \\
(47) + (48) + (52) &= 0, \\
(65) + (66) - \frac{1}{2}(68) &= 0; \tag{A1}
\end{aligned}$$

$$\begin{aligned}
(1') + 3(4') &= 0, \\
(2') + (5') - (6') &= 0, \\
(7') + (14') + (20') &= 0, \\
(8') + (13') + (21') &= 0, \\
(9') + (15') + (19') &= 0, \\
(10') - (16') &= 0. \tag{A2}
\end{aligned}$$

Note that some of these relations neglect contributions from lower levels in the hierarchy.

APPENDIX B: RELATIONS INDUCED BY EOM

$$\begin{aligned}
(1) &= (4), \\
(11) &= (53) - \frac{2}{N_f}(67), \\
(15) &= -(61) - \frac{1}{N_f}(67), \\
(21) &= (36) - \frac{2}{N_f}(66), \\
(49) &= (70) - \frac{1}{N_f}(73), \\
(51) &= (71) - \frac{1}{N_f}(74); \tag{B1}
\end{aligned}$$

$$\begin{aligned}
(5') &= (25'), \\
(19') &= (26'). \tag{B2}
\end{aligned}$$

APPENDIX C: EPSILON RELATIONS

$$\begin{aligned}
(11) + 2(12) - (13) + (14) &= 0, \\
(15) - 2(16) - (17) &= 0, \\
(24) - (25) &= 0, \\
2(24) + (26) - (27) &= 0, \\
(38) - 2(40) + (42) &= 0, \\
(44) - 2(46) + (48) &= 0, \\
(49) - 2(50) &= 0, \\
(51) - 2(52) &= 0, \\
2(55) + (56) - (57) &= 0, \\
2(55) - (58) + (59) &= 0, \\
2(56) + 2(58) + (60) &= 0, \\
(62) - 2(63) &= 0, \\
(62) + (64) &= 0, \\
(72) &= 0; \tag{C1}
\end{aligned}$$

$$\begin{aligned}
2(1') + 3(2') &= 0, \\
(1') + 3(3') &= 0, \\
2(7') + 2(8') + (9') &= 0, \\
2(13') + 2(14') + (15') &= 0, \\
(13') + 2(16') + (17') &= 0, \\
(14') - 2(16') - (18') &= 0,
\end{aligned}$$

$$\begin{aligned}
2(22') + (23') &= 0, \\
2(22') - (24') &= 0, \\
(27') - (29') &= 0, \\
(28') + (29') &= 0.
\end{aligned} \tag{C2}$$

APPENDIX D: RELATIONS INDUCED BY BIANCHI IDENTITIES

$$\begin{aligned}
(10) - \frac{1}{4}(26) &= 0, \\
(32) - \frac{1}{4}(33) &= 0, \\
(35) + \frac{1}{4}(54) &= 0, \\
(36) - \frac{1}{4}(53) + \frac{1}{2}(61) &= 0, \\
(42) + \frac{1}{4}(57) + \frac{1}{4}(62) &= 0, \\
(48) - \frac{1}{4}(58) - \frac{1}{4}(64) &= 0, \\
(66) - \frac{1}{2}(67) &= 0;
\end{aligned} \tag{D1}$$

$$\begin{aligned}
(8') - 2(10') &= 0, \\
(9') - 2(11') &= 0, \\
(12') &= 0, \\
(18') + \frac{1}{4}(23') &= 0.
\end{aligned} \tag{D2}$$

APPENDIX E: TRACE RELATIONS

$SU(2)$:

$$\begin{aligned}
(2) &= 0, \\
(4) &= 0, \\
(12) &= 0, \\
(13) &= 0, \\
(24) &= 0, \\
(26) &= 0, \\
(30) &= 0, \\
(31) &= 0, \\
(37) &= 0, \\
(43) &= 0,
\end{aligned}$$

$$\begin{aligned}
(53) + 2(61) &= 0, \\
(53) - (67) &= 0, \\
(54) + 2(68) &= 0, \\
(58) &= 0, \\
(59) &= 0, \\
(60) &= 0, \\
(63) &= 0, \\
2(70) - (73) &= 0, \\
2(71) - (74) &= 0;
\end{aligned} \tag{E1}$$

$SU(3)$:

$$(26) + (27) + 2(30) + (31) = 0. \tag{E2}$$

REFERENCES

- [1] S. Weinberg, *Physica* **96A**, 327 (1979).
- [2] J. Gasser and H. Leutwyler, *Ann. Phys.* **158**, 142 (1984).
- [3] J. Gasser and H. Leutwyler, *Nucl. Phys.* **B250**, 465 (1985).
- [4] A. M. Bernstein, D. Drechsel, and Th. Walcher (Eds.), *Chiral Dynamics: Theory and Experiment*, Proceedings, Mainz, Germany, 1997 (Springer, Berlin, 1998).
- [5] Y. Nambu, *Phys. Rev. Lett.* **4**, 380 (1960).
- [6] J. Goldstone, *Nuovo Cim.* **19**, 154 (1961).
- [7] J. Goldstone, A. Salam, and S. Weinberg, *Phys. Rev.* **127**, 965 (1962).
- [8] J. Stern, *Light Quark Masses and Condensates in QCD*, in Ref. [4].
- [9] H. W. Fearing and S. Scherer, *Phys. Rev. D* **53**, 315 (1996).
- [10] J. Bijnens, G. Colangelo, and G. Ecker, *JHEP* **9902**, 20 (1999).
- [11] R. Akhouri and A. Alfakih, *Ann. Phys. (N.Y.)* **210**, 81 (1991).
- [12] L. H. Ryder, *Quantum Field Theory* (Cambridge University Press, Cambridge, England, 1986), Chap. 3.6.
- [13] S. Weinberg, *The Quantum Theory of Fields, Vol. II* (Cambridge University Press, Cambridge, England, 1996), Chap. 15.3.
- [14] D. Issler, preprint SLAC-PUB-4943, 1990 (unpublished).